

Elite Hi-K

12-10-20 +

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N).....	12.00 %
<i>3% total N from stabilized Nitrogen</i>	
Available Phosphoric Acid (P ₂ O ₅).....	10.00 %
Soluble Potash (K ₂ O).....	20.00 %
Sulphur (S).....	12.00 %
Magnesium (Mg)	3.00 %
Boron (B).....	0.20 %
Zinc (Zn).....	0.10 %



FEATURES

- 25% of total N from stabilized nitrogen

BENEFITS:

Designed for use in either blueberries or raspberries where soils test low in potassium.

DIRECTIONS:

Rates may vary depending on soil types, crop history and variety. The following rates are correct for most plantings on organic soils in the Fraser Valley.

APPLICATION RATES:

Raspberries: *Meeker, Willamette:* 2 applications of 300 lbs per acre. *Skeena, Chilcotin, Chilliwack, Haida:* 2 applications of 250 lbs per acre. *Comox, Tulameen:* 300 lbs/acre sidedressed.

Blueberries: *1 - 3 Years:* Use 15-8-11 (Premium Berry Blend). *3+ Years:* Use 12-10-20+micros at 2 oz per plant per year of age to a maximum of 16 oz per plant. This application should be split; half in April and the remaining half in mid-May. On a standard 4 x 10 ft spacing, this amounts to 75 lbs/acre per year(s) of planting applied twice. Example: 3 year plants, 3 x 75 = 225 lbs/acre applied twice. Place fertilizer at the drip line of the plant (approximately outside edge of foliage).

Strawberries, all varieties, at planting: 300 lbs/acre sidedressed

Currants, Gooseberries, all varieties: 1 - 2 applications of 200 lbs/acre

As none of the nitrogen is coated in this blend, it is best banded into the soil to limit volatilization to the air. It can be topdressed where rain is forecast.

It is strongly recommended that the producer conduct tissue and soil tests. Fertilizer applications should be adjusted accordingly to avoid excessive build-up or deficiencies of one or more nutrients.

CAUTION:

Avoid contact with eyes and prolonged contact with skin. If eyes or skin are exposed, flush with water. Do not use on container stock or in greenhouse production. This is not a label; these guidelines are based on the experiences of producers in the Fraser Valley. Always refer to soil test results before use.